

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11 Publication number:

0 072 450
A1

12

EUROPEAN PATENT APPLICATION

21 Application number: 82106522.4

51 Int. Cl.³: C 12 Q 1/54
C 12 Q 1/28

22 Date of filing: 20.07.82

30 Priority: 03.08.81 US 289455

43 Date of publication of application:
23.02.83 Bulletin 83/8

84 Designated Contracting States:
DE FR GB

71 Applicant: MILES LABORATORIES INC.
1127 Myrtle Street
Elkhart Indiana 46514(US)

72 Inventor: Tybach, Richard J.
50966 George Drive
Elkhart Indiana 46514(US)

74 Representative: Senftl, Hannes, Dr. et al,
c/o Bayer AG Zentralbereich Patente Marken und
Lizenzen
D-5090 Leverkusen 1, Bayerwerk(DE)

54 Test device for lactase activity in a meconium sample.

57 A test device for the detection of lactase activity in a meconium sample is disclosed. The test device is a carrier matrix which has lactose, a glucose assay system and a nonionic detergent incorporated therein.

EP 0 072 450 A1

TEST DEVICE FOR LACTASE ACTIVITY
IN A MECONIUM SAMPLE

BACKGROUND OF THE INVENTION
AND PRIOR ART

5 Cystic fibrosis (CF) is a genetic illness which
has as its clinical features chronic pulmonary disease,
pancreatic dysfunction and a high concentration of
sodium and chloride in the sweat. Mucous plugs the
pancreatic ducts, the intestinal mucous glands and the
10 bronchial tree.

Studies have demonstrated raised levels of pro-
tein, predominately albumin, in meconium from infants
with CF [See: Pediatrics, 21:635-641 (1958)]. A test
device for the detection of albumin in meconium greater
15 than 20 milligrams per gram (mg/g) dry weight of
meconium has been developed. [See Pediatrics, 55:35-38
(1975)]. Such test devices suffer from a failure to
detect all newborns with CF, i.e., low sensitivity,
and an unacceptable number of false-positives, i.e.,
20 low specificity. Because of the emotional stress
caused by a false-positive test result for CF, re-
duction in the number of false-positive test results
in a CF test is desirable.

Pediatrics, 56:782-787 (1975) reported an increase
25 in disaccharidase activities in meconium from infants
with CF and suggested measurement of lactase activity

- 2 -

as well as the other activities, as a CF diagnostic test. Am. J. Dis. Child., 132:1112-1114 (1978) describes a study which included an assay test for lactase and β -D-fucosidase activity in conjunction
5 with an assay for albumin, in meconium. The authors suggest that the addition of the lactase and fucosidase assay would reduce the occurrence of false-positive test results.

The lactase activity test described above in-
10 volved a liquid assay test. The meconium sample was placed in a test tube and a 3 percent lactose solution in a maleate buffer added. After incubation for a period of 15 minutes, lactase activity was detected by the presence of glucose, demonstrated by immersing a
15 glucose reagent strip into the mixture and after 10 minutes, examining the glucose reagent strip for a blue color (positive test).

Use of a liquid assay test system for lactase activity detection in meconium samples is inconvenient.
20 Because the meconium sample is a tarry-like mass, the sample must be first homogenized. As described in Am. J. Dis. Child., *supra*, the sample is homogenized with vigorous agitation in a buffer solution. Other workers have resorted to ultrasonic techniques to obtain
25 homogenization. In order to properly prepare the sample for testing, the sample must be incubated for a period of from 15 minutes to one hour. In addition, the liquid assay test system requires the preparation of solutions for individual tests. In order to
30 encourage greater screening for CF in infants, a more convenient test system is required, preferably a dip-and-read reagent strip. The present invention is directed to such a dip-and-read test device.

SUMMARY OF THE INVENTION

The present invention is directed to a device for detecting lactase activity in a meconium sample. The device is a carrier matrix impregnated with lactose, 5 a glucose assay system, and a nonionic detergent.

DETAILED DESCRIPTION OF THE INVENTION

The test device can be prepared as elongated sheets of carrier material which have been incorporated with lactose, a glucose assay system and a 10 nonionic detergent. These elongated sheets may take the form of bulk rolls, such as of filter paper material. The device is prepared by incorporating a carrier with a solution containing lactose, a glucose assay system and a nonionic detergent, and thereafter 15 drying the impregnated carrier. The term carrier refers to matrices which are insoluble in and maintain their structural integrity when exposed to physiological or other liquids. Suitable matrices which can be used include paper, cellulose, wood, synthetic resin 20 fleeces, glass fiber, nonwoven and woven fabrics, gelatin, various organic polymers, such as polypropylene, and other organic materials well known as film formers to those skilled in the art. Alternatively, the carrier may take the form of a pressed or molded 25 tablet containing conventional carrier material. For convenience, the carrier can be suitably attached to an insoluble support or handle member which can be made from polystyrene. Incorporation of the carrier with the lactase-detecting solution can be effected by 30 suitable techniques, such as by impregnating, printing or spraying the test composition onto the carrier.

- 4 -

From about 1 to 4 percent by weight of lactose is present. A preferred amount is about 3 percent by weight. The glucose assay system includes a 3,3',5,5'-tetraalkylbenzidine indicator, wherein alkyl is a
5 C₁-C₄ alkyl; 3,3',5,5'-tetramethylbenzidine is preferred. Others which can also be used include 3-methyl, 3'-methyl, 5-ethyl, 5'-ethyl benzidine and 3,3',5,5'-tetraethylbenzidine.

The glucose assay system includes glucose enzymes
10 which will react with glucose, produced by conversion of lactose to glucose by lactase present in meconium to produce a predetermined reaction product, such as hydrogen peroxide. For example, glucose oxidase obtained from molds can be used.

15 Preferably, a dual enzyme system is present: one enzyme transforms glucose to produce hydrogen peroxide, whereas the other enzyme has peroxidative activity. Substances having peroxidative activity which are useful in the present invention can be chosen from
20 various organic and inorganic sources. Plant peroxidases, such as horseradish peroxidase or potato peroxidase, can be used. Inorganic compounds having peroxidative activity include iodides, such as sodium and ammonium iodides, and molybdates, such as potas-
25 sium and ammonium molybdates. In addition, urohemin and a number of other porphyrin substances having peroxidative activity can be used. Other substances which are not enzymes, but which have peroxidative activity include such compounds as iron sulfocyanate,
30 iron tannate, ferrous ferrocyanide, potassium chromic sulfate and the like.

Other glucose assay systems known to those skilled in the art are usable in the present invention. For example, those using glucose dehydrogenase, which
35 produce color in the presence of tetrazolium salts.

- 5 -

The detergent present in the composition is a nonionic detergent. The use of such a detergent in the formulation described enables the production of a test device which has increased sensitivity, e.g., in the presence of a lactase-containing meconium sample, the strip develops a positive color within 1 to 2 minutes. Nonionic detergents suitable for use in the present invention are alkanolamides, ethoxy alkanolamides, ethoxy phenols and ethoxy fatty alcohols.

10 The amount of nonionic detergent used can be in the range from about 0.1 to 1.0 percent by weight (w/v). A preferred amount is about 0.25 percent by weight. The presence of a greater amount of detergent may be deleterious to enzymes present in the test
15 device.

The test device of the present invention has a high sensitivity for detection of lactase activity. The meconium sample can be tested with the test device without dipping in water. It is only necessary to
20 smear a very thin film of meconium on the test device. If the sample has lactase activity, an easily visible blue color develops directly beneath the meconium.

An interpolymer of methylvinyl ether and maleic anhydride is also useful in the formulation of the
25 lactase detecting test device of the present invention. One such interpolymer is marketed commercially as Gantrez AN-139 by GAF New York, New York. When this interpolymer is dissolved in an alcohol it forms a partial ester derivative, and when the interpolymer is
30 dissolved in water it forms an acid derivative. Since test means prepared in accordance with the present invention are typically prepared from aqueous alcohol solutions, test compositions in the final product will contain either an acid derivative or a partial ester
35 derivative or a mixture of said derivatives. The

presence of the above described interpolymer derivatives along with polyvinyl pyrrolidone (PVP) having, for example, an average molecular weight of about 40,000, greatly enhances the color formed when color forming indicators are oxidized by hydrogen peroxide in the presence of peroxidase. This enhancement of color aids in detecting the presence of glucose produced by the conversion of lactose to glucose, and hence a positive indication of lactase activity, in the meconium sample.

Horseradish peroxidase and glucose oxidase used in the example were obtained from the Research Products Division, Miles Laboratories, Inc., Elkhart, Indiana. A copolymer of methyl vinyl ether and maleic anhydride (Gantrez AN-139) and polyvinyl pyrrolidone (PVP) were obtained from GAF. The solvent used in preparing the solutions can be water, physiological solutions, organic solvents, such as methanol, or mixtures thereof.

20

EXAMPLE 1

Dip-and-read test devices for the detection of lactase activity were prepared using a two-step impregnating procedure. A first solution was prepared by mixing together the substances shown in Table 1 below.

- 7 -

TABLE 1

	<u>Component</u>	<u>Amount</u>	<u>Final Concentration</u>
			<u>in Dip</u>
5	(w/v) polyvinylpyrrol- idone (PVP)*	30.0 milli- liter (ml)	3.0 %
	1.37 M Citrate Buffer, pH 5.0	15.0 ml	0.20 %
10	poly(methylvinyl ether maleic anhydride) 10% (w/v)	7.5 ml	0.75 %
	polyethoxy oleyl** alcohol 10% (w/v)	2.5 ml	0.25 %
15	Glucose Oxidase (4878 IU/ml)	1.8 ml	87.8 inter- national units per milliliter (IU/ml)
	Peroxidase (153 U/mg)	50.0 milli- grams (mg)	0.5 mg/ml
20	Lactose	3.0 grams (g)	3.0 %
	20% w/v Ascorbic acid***	10 micro- liter (ul)	0.002 %
	Distilled Water	to bring volume to 100 ml	

* A PVP commercially available from GAF under the
25 trade designation K29-32.

** A polyethoxylated fatty alcohol commercially avail-
able from GAF under the trade designation ON 870.

30 ***Added to reduce oxidized indicator which forms as
a result of peroxide contaminants in the PVP. Amount
to add determined by titration of the dip to the
equivalence point.

- 8 -

A sheet of Whatman 54 filter paper was impregnated to saturation with this impregnating solution and dried at 50-55°C. for about 15 minutes. The impregnated sheet was then impregnated to saturation with a 5 millimolar (mM) solution of 3,3',5,5'-tetramethylbenzidine in acetone containing 0.1 percent ON-870, and dried at 50-55°C. for one minute.

The paper so prepared was cut to 0.2 centimeter (cm) x 0.4 cm to form test devices. The devices were then backed with double-faced adhesive tape and fixed thereby to polystyrene support members. Test devices were evaluated in two formats. In the first, a thin-layer of meconium was smeared on the test device. Care was taken to remove all excess by scraping the test device with a spatula. A pathological meconium sample, known to have high lactase activity showed blue color development where it was in contact with the test device, within ten seconds. Apparently, enough water is present in the meconium for the enzymatic reaction to occur in the test device.

In the second format, a drop of water was placed on the test device after application of the meconium. This served to enhance color development.

As a control, a meconium sample, known to have no lactase activity was tested with the test device of the present invention. The test was negative for lactase activity.

Accordingly, the tests of this example showed that the test device of the invention is capable of detecting lactase activity in a meconium sample accurately, quickly and easily.

WHAT IS CLAIMED IS:

1. A test device for the detection of lactase in a meconium sample which comprises a carrier matrix, and incorporated therein, lactose, a glucose
5 assay system and a nonionic detergent.
2. A test device as claimed in Claim 1 wherein the glucose assay system comprises a 3,3',5,5'-tetraalkylbenzidine indicator, glucose oxidase and a peroxidatively active substance.
- 10 3. A test device as claimed in Claim 1 wherein the peroxidatively active substance is peroxidase.
4. A test device as claimed in Claim 1 wherein the indicator is 3,3',5,5'-tetramethylbenzidine.
5. A test device as claimed in Claim 1 wherein from
15 about 1 to 4 percent by weight of lactose is present.
6. A test device as claimed in Claim 1 wherein from about 0.1 to 1 percent by weight of nonionic detergent is present.
- 20 7. A test device as claimed in Claim 1 wherein the nonionic detergent is a polyethoxy fatty alcohol.
8. A test device for detection of lactase activity in a meconium sample which comprises a carrier matrix, and incorporated therein, lactose,
25 3,3',5,5'-tetramethylbenzidine, glucose oxidase, peroxidase, and polyethoxy oleyl alcohol.

9. A process for detecting lactase activity in a meconium sample which comprises the steps of contacting said sample with a test device comprising a carrier matrix impregnated with lactose, a glucose assay system and a
5 nonionic detergent and observing a detectable response produced by said test device.
10. A process as claimed in claim 9 wherein the glucose assay system comprises a 3,3', 5,5'-tetraalkylbenzidine indi-
10 cator, glucose oxidase and a peroxidatively active substance.
11. A process for producing a test device for detecting lactase activity by impregnating a carrier with a composition which includes lactose, a glucose assay system and a nonionic detergent and drying the carrier.
- 15 12. A process as claimed in claim 11 wherein the glucose assay system comprises a 3,3', 5,5'-tetraalkylbenzidine indicator, glucose oxidase and a peroxidatively active substance.



European Patent
Office

EUROPEAN SEARCH REPORT

0072450
Application number

EP 82 10 6522

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
Y	--- GB-A-1 445 793 (ALEA-LAVAL AB) *The whole document*	1-12	C 12 Q 1/54 C 12 Q 1/28
X,Y	--- EP-A-0 029 917 (MILES LABORATORIES INC.) *Example 1, line 12; claims 3,4*	1-12	
X,Y	--- US-A-4 273 868 (BERT WALTER) *Abstract; claim 1*	1-12	
A	--- CHEMICAL ABSTRACTS, vol. 94, no. 17, April 1981, page 384, no. 135467u, Columbus Ohio (USA); G.MACHILL et al.: "Proposals to the pharmacopeia of GDR, 2nd edition. Diagnostic laboratory methods. Detection of mucoviscidosis". & ZENTRALBL. PHARM., PHARMAKOTHER. LABORATORIUMSDIAGN. 1980, 119(12), 1321-5. *Abstract*		
			TECHNICAL FIELDS SEARCHED (Int. Cl. 3)
			C 12 Q G 01 N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 15-11-1982	Examiner OSBORNE H.H.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	